



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.


JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

IV. "On some new Colouring Matters." By ARTHUR H. CHURCH and WILLIAM H. PERKIN. Communicated by A. W. HOFMANN, Ph.D., F.R.S. Received February 5, 1856.


Nascent hydrogen, acting upon an alcoholic solution of dinitrobenzole or of nitraniline, produces a crimson coloration, due to the formation of a new substance, to which we have given the name *nitrosophenyline*.

This new body presents some remarkable properties. It fuses below  $100^{\circ}\text{C}$ .; is uncrystallizable, and not volatile without decomposition; it dissolves in alcohol with an orange-red tint, and an alcoholic solution containing only  $\cdot 2$  per cent., although perfectly transparent to transmitted light, presents a flame-coloured luminous opacity in reflected light. Nitrosophenyline dissolves in hydrochloric acid, producing an intense crimson colour, which is changed to a yellowish-brown by alkalies and is restored by acids.

The analysis of nitrosophenyline has led us to the formula



which may be written thus:— $\text{C}_{12}\overset{\text{H}_6}{\text{NO}_2}\text{N}$ , and so may be viewed as aniline, in which 1 equiv. of hydrogen is replaced by 1 equiv. of binoxide of nitrogen: the following equation sufficiently explains the formation of nitrosophenyline:—




We have produced from all the dinitro-compounds we have yet experimented upon, colouring matters similar to nitrosophenyline: the following is a list of such dinitro-compounds:—

1. Dinitrobenzole . . . .  $\text{C}_{12}\text{H}_4\text{2NO}_4$ .
2. Dinitrotoluole. . . . .  $\text{C}_{14}\text{H}_6\text{2NO}_4$ .
3. Dinitroxylol . . . . .  $\text{C}_{16}\text{H}_8\text{2NO}_4$ .
4. Dinitroxylol . . . . .  $\text{C}_{18}\text{H}_{10}\text{2NO}_4$ .
5. Dinitrocymole. . . . .  $\text{C}_{20}\text{H}_{12}\text{2NO}_4$ .
6. Dinitronaphthaline. .  $\text{C}_{20}\text{H}_6\text{2NO}_4$ .

We have examined minutely the colouring substance produced in

the case of dinitronaphthalene. It proves to be perfectly analogous in composition with nitrosophenylene; in properties also it is similar; and from its alcoholic solution it may be obtained in crystals, having a lustre somewhat similar to that of murexide: its formula, as deduced from our analysis, is



which we may arrange thus:— $\text{C}_{20} \text{H}_8 \text{NO}_2$ , and so view it as naphthylamine in which 1 equiv. of hydrogen has been replaced by 1 equiv. of binoxide of nitrogen. This substance we term nitrosophenylene. It may likewise be obtained by the action of nitrous acid on naphthylamine, or of nitrite of potassium upon the hydrochlorate of naphthylamine: the following equations represent the three processes for its formation:—

1.  $\text{C}_{20} \text{H}_6 2\text{NO}_4 + 8\text{H} = \text{C}_{20} \text{H}_8 \text{N}_2 \text{O}_2 + 6\text{HO}.$
2.  $\text{C}_{20} \text{H}_9 \text{N} + \text{NO}_3 = \text{C}_{20} \text{H}_8 \text{N}_2 \text{O}_2 + \text{HO}.$
3.  $\text{C}_{20} \text{H}_{10} \text{N, Cl} + \text{KNO}_4 = \text{C}_{20} \text{H}_8 \text{N}_2 \text{O}_2 + 2\text{HO} + \text{KCl}.$

*February 28, 1856.*

The LORD WROTTESLEY, President, in the Chair.

The following communications were read:—

- I. The following Letter was read, from Professor HANSTEEN of Christiania, For. Mem. R.S.:—

*To the Royal Society of London.*

As a Corresponding Member of the Royal Society, I have the honour herewith to transmit a Research “On the Secular Changes of the Magnetical System of the Earth, and more specially on the Secular Variation of the Magnetical Inclination in the Northern Temperate Zone,” separately printed from the ‘Memoirs of the Roy. Soc. of Sciences of Copenhagen.’ By calculating newer and more ancient observations of the magnetical declination, I have ascertained the movement of the four magnetical polar regions, which I had